
Little brown myotis maternity roost surveys: Copper River Basin, 2016

Jesika P. Reimer, Laura A. Kaupas and Paul A. Schuette



October 2016

Prepared by

Alaska Center for Conservation
Science University of Alaska
Anchorage
Beatrice MacDonald Hall
3211 Providence Drive
Anchorage, Alaska 99508

Prepared in cooperation with

Threatened, Endangered and Diversity
Program Alaska Department of Fish
and Game
1255 W. 8th Street
Juneau, Alaska 99811



Recommended citation:

Reimer, J.P., L.A. Kaupas and P.A. Schuette. 2016. Little brown myotis maternity roost surveys: Copper River basin, 2016. [Report] Alaska Center for Conservation Science, University of Alaska Anchorage.

Table of Contents

| | |
|---|-----|
| Introduction | 1 |
| Methods..... | 1 |
| <i>Emergence counts and estimated colony size</i> | 1 |
| <i>Capture surveys</i> | 2 |
| Results..... | 2 |
| Discussion..... | 5 |
| Ongoing Research | 6 |
| Acknowledgements..... | 6 |
| Literature cited..... | 7 |
| APPENDIX A – Maternity roost profiles | A-i |
| Roost ID: RM-16-01 | A-1 |
| Roost ID: RM-16-02 | A-2 |
| Roost ID: RM-16-03 | A-3 |
| Roost ID: RM-16-04 | A-4 |
| Roost ID: RM-16-05 | A-5 |
| Roost ID: RM-16-06 | A-6 |
| APPENDIX B – Artificial Roost Pilot Project..... | B-i |
| Artificial Roost ID: RB-16-01 | B-1 |
| Artificial Roost ID: RB-16-02 | B-2 |
| Artificial Roost ID: RB-16-03 | B-3 |
| APPENDIX C – Banding Records - 2016 | C-1 |

Figures and Tables

| | |
|--|-----|
| Figure 1: Locations of <i>Myotis lucifugus</i> maternity colonies surveyed during July, 2016..... | 2 |
| Figure 2: Age class and reproductive status capture profile of <i>Myotis lucifugus</i> captured during survey events in Alaska, July 2016. | 4 |
| Figure 3: Little brown myotis (<i>Myotis luficugus</i>) maternity roost RM-16-01 | A-1 |
| Figure 4: Little brown myotis (<i>Myotis luficugus</i>) maternity roost RM-16-02 | A-2 |
| Figure 5: Little brown myotis (<i>Myotis luficugus</i>) maternity roost RM-16-03 | A-3 |
| Figure 6: Little brown myotis (<i>Myotis luficugus</i>) maternity roost RM-16-04 | A-4 |
| Figure 7: Little brown myotis (<i>Myotis luficugus</i>) maternity roost RM-16-05 | A-5 |
| Figure 8: Little brown myotis (<i>Myotis luficugus</i>) maternity roost RM-16-06 | A-6 |
| Figure 9: Installation of artificial bat maternity roost RB-16-01 (style: rocket box) behind an active <i>Myotis lucifugus</i> maternity roost (RM-16-03). | B-1 |
| Figure 10: Construction of artificial bat roost RB-16-02 (style: rocket box). | B-2 |
| Figure 11: Installation of artificial bat maternity roost RB-16-03 (style: rocket box) behind an active <i>Myotis lucifugus</i> maternity roost (RM-16-05; not in picture). | B-3 |
| Table 1: Location and surrounding habitat of <i>Myotis lucifugus</i> maternity roosts along the Copper River, Alaska. | 3 |
| Table 2: Structure characteristics and estimated colony size of <i>Myotis lucifugus</i> maternity roosts along the Copper River, Alaska. | 3 |
| Table 3: Reproduction rates of adult female <i>Myotis lucifugus</i> observed at maternity roosts during July 2016 along the Copper River basin, Alaska. | 3 |
| Table 4: Prevalence of ear tissue damage observed at <i>Myotis lucifugus</i> maternity colonies along the Copper River delta, AK, surveyed during July 2016..... | 4 |
| Table 5: Little brown myotis (<i>Myotis lucifugus</i>) banding records for captures during July 2016 surveys.C-1 | |

Introduction

Little brown myotis are widespread across North America, and reach their northernmost extent in Alaska, Yukon and the Northwest Territories, Canada (Wilson et al. 2014). While bat research has been ongoing throughout Alaska since the early 1990's (Parker et al. 1996), relatively little is known about population size or trends in interior Alaska. In addition, while bats in southeast Alaska have been observed hibernating in caves, scree fields and root wads (K. Blejwas, pers. comm. 2016; Parker et al. 1996), winter locations for bats that summer in interior Alaska are still unknown.

Since 2007, little brown myotis have been dying by the millions in eastern North America due to white-nose syndrome (WNS), raising the alarm across their range. WNS is a disease that has a 90% mortality rate for infected little brown myotis colonies (USFWS 2016). It is currently present in 27 states and 5 Canadian provinces in eastern North America, and has killed over 6 million bats to date. As of 31 March 2016, white-nose syndrome was confirmed in Washington (USGS 2016), over 1300 miles from the nearest infection site. It has not yet arrived in Alaska, but if/when it does, having an understanding of species distribution, seasonal activity and abundance will assist in detecting the arrival and impact of WNS and enhance our ability to take appropriate action. Since hibernation behavior and over-wintering sites for little brown myotis are still unknown for bats in interior Alaska, testing for WNS at winter sites is not yet possible. However, given the site fidelity shown by females to summer maternity colonies, investigating and monitoring population fluctuations at these congregation sites may be the next best alternative (Dobony et al. 2011).

Our goal is to identify a network of little brown myotis (*Myotis lucifugus*) maternity colonies throughout Alaska to assist with studying and monitoring bat populations over time. In particular we plan to: a) begin establishing baseline population levels at maternity sites to effectively monitor fluctuations indicating the potential arrival of WNS to Alaska; b) investigate population dynamics inclusive of estimating colony size, survival, and reproductive rates, which will help inform management decisions; and c) build a network of citizen scientists to support monitoring efforts at maternity colonies throughout Alaska.

Methods

During the first year of our study (2016) we focused on six maternity colonies in local residences, including five along the Copper River Valley and one north of Anchorage in Wasilla. We spent two nights at each roost site counting emergence numbers and mist-netting to survey individuals in the colony. In addition, we recorded roost measurements including overall size, material type and general location.

Emergence counts and estimated colony size

Emergence counts were performed at each roost location on the first night of arrival at the site. Two to four observers were situated around the roost twenty minutes before sun set. Each observer was assigned to watch a portion of the roost building, and tallied bats emerging using a handheld tally clicker. Observers continued to count emerging bats for a minimum of one hour after emergence commenced, and ceased counting after a fifteen minute period of inactivity at the roost, or if more than ten bats were seen re-entering the roost. Emergence counts were not performed during nights of inclement weather such as frequent or heavy rain or wind. For colonies where juveniles had not yet fledged (no juveniles were captured during the survey night), colony size was estimated to be the total emergence count. For colonies where fledged juvenile were captured during the survey night, colony

size was estimated by multiplying the total emergence count by the proportion of the survey population identified as adults (e.g. a roost with an emergence count of 100 and a capture survey consisting of 5% juveniles would result in an estimated colony size of 95 adults).

Capture surveys

During night two at each colony, standard capture techniques using mist nets were used to live-capture bats exiting the roost. Measurements were recorded for each capture including ear length, forearm length, mass, sex, tooth wear (indication of age), and reproductive status. Each bat was fitted with a metal lipped band engraved with a unique identifier for future identification during reproductive and population monitoring studies (see Appendix C). All capture and bat handling techniques strictly adhered to the National White-Nose Syndrome Decontamination Protocol (Version 04.12.2016). All animal handling was performed under IACUC Permit # 918743-1.

Results

We performed roost emergence counts and mist-netting surveys of little brown myotis at six known maternity roosts during July 2016 (Figure 1). Roosts were located in a variety of buildings including private residences, recreational cabins, abandoned cabins, and a large barn. All buildings were constructed of wood and majority of the buildings were fitted with a tin roof, with the exception of roost RM-16-02 which had a wood roof covered with tarpaper (see Appendix A). Majority of the roosts were within 200 meters of the nearest water body, and all roosts were within 20 m from the nearest forest edge (Table 1).




Figure 1: Locations of *Myotis lucifugus* maternity colonies surveyed during July, 2016

Table 1: Location and surrounding habitat of *Myotis lucifugus* maternity roosts along the Copper River, Alaska.

| Roost | Latitude (N) | Longitude (W) | Distance from forest (m) | Distance from water (m) | Nearest waterbody (type) |
|----------|--------------|---------------|--------------------------|-------------------------|--------------------------|
| RM-16-01 | 61.50 | -149.59 | 3 | 20 | River |
| RM-16-02 | 62.89 | -143.68 | 20 | 100 | Beaver pond |
| RM-16-03 | 62.69 | -143.82 | 15 | 100 | Small ponds, streams |
| RM-16-04 | 62.30 | -145.31 | 20 | 200 | River |
| RM-16-05 | 61.70 | -144.86 | 20 | > 500 | unknown |
| RM-16-06 | 61.68 | -144.69 | 10 | 500 | River |

Colony size ranged from approximately 50 to 470 adults and was not correlated with roost size (Table 2; Linear regression, $F_{1,4} = 0.53$, $p > 0.05$).

Table 2: Structure characteristics and estimated colony size of *Myotis lucifugus* maternity roosts along the Copper River, Alaska.

| Roost | Length (m) | Width (m) | Height (m) | Total Area (m ²) | Total Volume (m ³) | Roof Type | Estimated Colony size |
|----------|------------|-----------|------------|------------------------------|--------------------------------|-----------------------------|-----------------------|
| RM-16-01 | 47 | 13 | 13 | 611 | 7,943 | Tin - open ceiling | 48 |
| RM-16-02 | 11 | 9 | 7 | 99 | 693 | Shingle roll - closed attic | 417 [†] |
| RM-16-03 | 4 | 4 | 4 | 16 | 64 | Tin - closed ceiling | 57 |
| RM-16-04 | 6 | 5 | 7 | 30 | 210 | Tin - closed ceiling | 468 |
| RM-16-05 | 20 | 7 | 8 | 140 | 1,120 | Tin - closed ceiling | 66 [†] |
| RM-16-06 | 10 | 6 | 9 | 60 | 540 | Tin - closed ceiling | 45 [†] |

[†] Emergence count adjusted to account for fledged juveniles

In total, we captured 140 individual bats and banded 136 individuals (see Appendix C). Average forearm length for adult females was 38.6 ± 0.96 mm (range: 36.2 to 40.9 mm; $n = 121$) and average mass was 8.0 ± 0.7 g for lactating and non-reproductive females (range: 6.0 to 10.5 g; $n = 114$), and 10.3 ± 1.0 g for pregnant females (range: 8.5 to 12.0 g; $n = 8$). The average female reproductive rate across colonies was 64.5% and ranged from 33% to 94% at each colony (Table 3).

Table 3: Reproduction rates of adult female *Myotis lucifugus* observed at maternity roosts during July 2016 along the Copper River, Alaska.

| Roost | Netting Date | Reproductive Rate | Sample size | Estimated Colony size |
|----------|--------------|-------------------|-------------|-----------------------|
| RM-16-01 | 4-Jul-16 | 0.33 | 3 | 48 |
| RM-16-02 | 9-Jul-16 | 0.89* | 44 | 417 [†] |
| RM-16-03 | 8-Jul-16 | 0.64 | 22 | 57 |
| RM-16-04 | 11-Jul-16 | 0.57 | 28 | 468 |
| RM-16-05 | 13-Jul-16 | 0.94* | 16 | 66 [†] |
| RM-16-06 | 20-Jul-16 | 0.50 | 8 | 45 [†] |

* Statistically higher than the overall mean reproductive rate

[†] Emergence count adjusted to account for fledged juveniles

Juveniles were captured at three of the six sites and the first volant juvenile was captured on 9 July, however, non-volant juveniles were also observed at the roost entrance that night. Juvenile captures ranged in size from 36.1 mm to 38.5 mm forearm length (mean: 37.5 ± 0.82 , $n=13$) and 4.75 g to 7.5 g mass (mean: 6.5 ± 0.70 , $n = 13$). The largest proportion of juveniles were captured at the latest surveyed roost on 20 July (Figure 2).




Figure 2: Age class and reproductive status capture profile of *Myotis lucifugus* captured during survey events in Alaska, July 2016.

During capture events, we noticed damaged ear tissue in 12% of the total number of captures ($n = 140$). Ear tissue damage was present in 0% to 24% of the captured individuals at each site (Table 4).

Table 4: Prevalence of ear tissue damage observed at *Myotis lucifugus* maternity colonies along the Copper River delta, AK, surveyed during July 2016.

| Location | Captures with frostbite | Sample size | % captures with frostbite | Estimated colony size | % Population surveyed |
|--------------|-------------------------|-------------|---------------------------|-----------------------|-----------------------|
| RM-16-01 | 0 | 6 | 0.00 | 48 | 12.50 |
| RM-16-02 | 1 | 48 | 2.08 | 417 [†] | 11.50 |
| RM-16-03 | 5 | 23 | 21.74 | 57 | 40.35 |
| RM-16-04 | 5 | 28 | 17.86 | 468 | 5.98 |
| RM-16-05 | 5 | 20 | 25.00 | 66 [†] | 30.30 |
| RM-16-06 | 1 | 15 | 6.67 | 45 [†] | 33.33 |
| TOTAL | 17 | 140 | 12.14 | 1101 | 12.72 |

[†] Emergence count adjusted to account for fledged juveniles

There was no significant difference between standardized mass (mass/forearm length; ANOVA, $df = 120$, $R^2 = 0.0005$, $F = 0.06$, $p > 0.05$) or reproductive rate (ANOVA, $df = 120$, $R^2 = 0.0152$, $F = 1.84$, $p > 0.05$) of adult females with and without ear tissue damage.

Discussion

During this first year of study, we identified and surveyed six little brown myotis maternity colonies in an effort to build a roost monitoring network across Alaska. All six colonies inhabited man-made roosts that varied in size, materials, human-occupancy, and nearby habitat (Appendix A). They were all within 20 meters from the nearest forest edge, and five of the six colonies had a water-body nearby (Table 1). This close proximity to the forest may be preferential for foraging and predation avoidance given the relatively bright nights at high latitudes (Rydell 1992). Colony population sizes ranged from 50 to nearly 500 adults and did not appear to be influenced by the size of the roost (Table 2). These man-made roost types and estimated colony sizes are consistent with little brown myotis roost selection and population numbers reported for colonies throughout their North American range (Anthony and Kunz 1977, Jung 2013, Randall et al. 2014, Smith 1940, Talerico 2008, Wilson *et al.* 2014).

Female reproductive rates varied across surveyed colonies with an average reproductive rate of 64.5% (Table 3). This is relatively low compared to maternity colonies farther south in places such as New Hampshire (87 to 99%; Frick *et al.* 2010) and the eastern US (> 96%; Cagle and Cockram 1943, Humphrey and Cope 1976). These lower reproductive rates are similar to those observed at other northern latitudes such as Yukon (33 to 74%; Talerico 2008) and the Northwest Territories (49 to 79%; Reimer 2013), Canada, and consistent with a documented trend of declining reproductive rates associated with increasing latitudes (Barclay et al. 2004). It should be noted however, that two maternity colonies surveyed this summer had reproductive rates significantly higher than the overall average (89% and 94%; Table 3) and were more consistent with rates observed farther south, illustrating the high variability observed across colonies within a given year. Future research investigating roost characteristics such as temperature profiles, and ambient seasonal temperatures may shed light on the causes of reduced reproductive rates within various roosts and colonies throughout Alaska. Reproductive rates of little brown myotis are known to be influenced by climate (Frick *et al.* 2010), and future research will investigate seasonal temperatures in conjunction with internal roost temperatures and reproductive rates across colonies.

While we did not do extensive temporal surveys at each colony to determine timing of parturition, lactation and fledging, our surveys at each site seem to suggest that overall, juveniles are fledging during the middle of July, which is consistent with what is seen at other northern sites (e.g. Northwest Territories; Reimer 2013). Interestingly, there appeared to be large variation in parturition dates both within and between colonies. At roost RM-16-02, pregnant females, lactating females, non-volant juveniles and fledged juveniles were all captured or observed (Figure 2). This overlap in reproductive stages has also been observed at northern sites in the Yukon (Talerico 2008) and may be linked to cool spring temperatures, variation in individual female fitness, and/or variation in torpor-use by individuals (Frick *et al.* 2010, Racey and Swift 1981).

During our capture surveys we noted a large number of bats exhibiting signs of ear tissue damage (Table 4). Currently in the literature there are only two papers that reference similar findings; LaVal and LaVal (1970) studied myotis species in Missouri and suggested that truncated ears were a sex-linked gene mutation that affected only males, while Kurta and Kweiscinski (2007) did histological studies of myotis ear tissue at hibernacula in Michigan and suggested that the truncated ears were the result of trauma most likely associated with frostbite. Since we observed damaged ear tissue in both males and females and the damage was not uniform across ears, we expect that the damage is similar to that reported by

Kurta and Kweiscinski (2007) and may be attributed to frostbite. Over the past few years, residents throughout Alaska have reported bats in human dwellings during the winter, and a recent radio tracking study during September suggested that bats may be spending the winter in buildings (Shively 2016). In addition, recent studies in southeast Alaska observed a small number of bats roosting in small cracks and crevices of scree fields and root wads (K. Blejwas, pers. comm. 2015). The high frequency of ear tissue damage observed in our study may support the idea that bats are spending the winter in roost sites throughout interior Alaska where temperatures fluctuate and drop to very low figures rather than migrating to more stable hibernacula farther south. While this winter-roosting behavior would be considered unusual compared to the rest of the species' range, winter roosting in buildings and scree fields has been observed for bat species in northern Norway (Michaelsen *et al.* 2013), and may be a strategy employed by bats at northern latitudes when relatively large, stable caves are lacking. Additional research to determine the definitive cause of the ear tissue damage, and the potential fitness effects is warranted.

In addition to monitoring existing maternity colonies, we worked with home owners to build artificial bat boxes in an effort to move bats out of human dwellings (Appendix B). We constructed three, single-chamber rockets boxes that were installed within close proximity to existing maternity colonies. Each bat box was outfitted with a hobo data logger and will be monitored for roost conditions and bat activity during 2017.

In conclusion, we consider this first summer of little brown myotis maternity colony surveys to have been a success in laying the groundwork for a long term monitoring program. We have identified wide variation in reproductive rates across colonies and reproductive phenology across individuals, and noted a high prevalence of damaged ear tissue amongst populations. Monitoring these six maternity colonies, with the help of local residents, will allow us to start following population trends at different sites to monitor for changes in population numbers at both a localized and region-wide level.

Ongoing Research

Data collection for the 2016 study period is still ongoing, and we currently have temperature loggers deployed in each of the six maternity roosts and each of the three artificial bat boxes. We will retrieve the temperature loggers during November 2016 and assess the temperature profiles of each roost in relation to ambient conditions. Future research plans include a second field season for 2017 during which we plan to add an additional six maternity roosts along the Tanana River to the roost monitoring network. We will also re-survey the six 2016 maternity roosts along the Copper River, to begin the second year of population data collection. Acoustic monitoring at these sites will commence in the spring (April) to identify when bats arrive at each maternity roost. If logistically feasible, acoustic monitoring will continue through the summer and autumn of 2017 to determine season length of female myotis across maternity colonies. Web-based outreach materials are currently being developed in collaboration with ADF&G to further engage citizen scientists in roost monitoring activities; and additional research into how environmental and roost conditions may impact reproductive rates and phenology is being considered.

Acknowledgements

This project was made possible through a co-operative funding agreement with the Alaska Department of Fish and Game - Threatened, Endangered and Diversity Program. We would like to thank the

numerous home owners who allowed us access to their homes and were eager to help us with our research. In particular we would like to thank: Herman Griesse, Joy and Steve of Huck Hobbit's, Ruby's family in Mentasta, Vonda and Robert of Gakona, Pete Peschang, and Ruth and Cliff McHenry. For assistance in the field, we thank M. Kohan, A. Neithercoat, B. Bernard and A. McKnown.

Literature cited

- Anthony, E.L.P. and T.H. Kunz. 1977. Feeding strategies of the little brown bat, *Myotis lucifugus*, in Southern New Hampshire. *Ecology* 58: 775-786.
- Barclay, R.M.R., J. Ulmer, C.J.A. MacKenzie, M.S. Thompson, L. Olson, J. McCool, E. Cropley and G. Poll. 2004. Variation in the reproductive rate of bats. *Canadian Journal of Zoology* 82: 688-693.
- Blejwas, K. pers. comm. 2015. Personal conversations with J. Reimer. September 2015. Regional Wildlife Biologist, Southeast Alaska (Region I), Threatened, Endangered and Diversity Program, Alaska Department of Fish and Game, Juneau, AK.
- Cagle, F.R. and L. Cockrum. 1943. Notes on a summer colony of *Myotis lucifugus lucifugus*. *Journal of Mammalogy* 24: 474-492
- Dobony, C A., A.C. Hicks, K.E. Langwig, R.I. von Linden, J.C. Okoniewski, and R.E. Rainbolt. 2011. Little brown myotis persist despite exposure to white-nose syndrome. *Journal of Fish and Wildlife Management*, 2: 190-195.
- Frick, W.F., D.S. Reynolds and T.H. Kunz. 2010. Influence of climate and reproductive timing on demography of little brown myotis *Myotis lucifugus*. *Journal of Animal Ecology* 79: 128-136
- Humphrey, S.R. and J.B. Cope. 1976. Population ecology of the little brown bat *Myotis lucifugus* in Indiana and south-central Kentucky. *American Society of Mammalogists. Special Publication* 4: 1-81.
- Jung, T.S. 2013. Estimating little brown bat (*Myotis lucifugus*) colony size in southern Yukon: a mark-recapture approach. Yukon Fish and Wildlife Branch Report TR-13-13. Whitehorse, Yukon, Canada.
- Michaelsen, T.C., O. Olsen and K.J. Grimstad. 2013. Roosts used by bats in late autumn and winter at northern latitudes in Norway. *Foila Zool.* 62: 297-303.
- Parker, D.I., J.A. Cook and S.W. Lewis. 1996. Effects of timber harvest on bat activity in Southeastern Alaska's temperate rainforests, pp. 227-292 in Barclay, R.M.R. and Brigham, R.M. (eds.). *Bats and Forests Symposium*. Research Branch, Ministry of Forests, Victoria BC, 23:1-292.
- Racey, P.A. and S.M. Swift. 1981. Variation in gestation length in a colony of pipistrelle bats (*Pipistrellus pipistrellus*) from year to year. *Journal of Reproduction and Fertility* 61:123-129.
- Randall, L.A., T.S. Jung and R.M.R. Barclay. 2014. Roost-site selection and movements of little brown myotis (*Myotis lucifugus*) in southwestern Yukon. *Northwestern Naturalist* 95:312-317.
- Rydell, J. 1992. Occurrence of bats in northernmost Sweden (65°N) and their feeding ecology in summer.
- Shively, R. 2016. Diet and habitat of the little brown bat (*Myotis lucifugus*) in interior and northern Alaska. [MS thesis] University of Alaska Fairbanks, 72 pp.
- Smith, R.W. 1940. The Land Mammals of Nova Scotia. *American Midland Naturalist* 24:213-241.

- Talerico, J. 2008. The behaviour, diet and morphology of little brown bats (*Myotis lucifugus*) near the northern extent of its range in Yukon, Canada. [MSc thesis] University of Calgary, 104 pp.
- U.S. Fish and Wildlife Service (USFWS). 2016. White-Nose Syndrome: North America's Response to the Devastating Bat Disease. Website: <https://www.whitenosesyndrome.org/faq/what-effect-white-nose-syndrome-bats>, [Accessed 3 March 2016].
- U.S. Geological Survey (USGS). 2016. News Release: Bat with white-nose syndrome confirmed in Washington State. Available online: <https://www.usgs.gov/news/bat-white-nose-syndrome-confirmed-washington-state>, [Accessed 2 March 2016].
- Wilson, J.M., J.P. Reimer, D. Allaire and C.L. Lausen. 2014. Diversity and distribution of bats in the Northwest Territories. *Northwestern Naturalist* 95:197-218.



APPENDIX A – Maternity roost profiles

Roost ID: RM-16-01

Species: *Myotis luficugus*, Little brown myotis

Date surveyed: 3 July 2016

Colony emergence count 2016: 48 individuals

 Bat exit/entrance point
 Flight pathway




Figure 3: Little brown myotis (*Myotis luficugus*) maternity roost RM-16-01

Roost size (length x width x height): 140 m x 40 m x 40 m

General location: Wasilla, AK


Comments: The roost is an old barn situated on private land. The roof is 'open', with bats roosting in the peak of the SW side. The nearest water source is a river approximately 20 meters from the SE side. The barn is currently being used as storage.

Roost ID: RM-16-02

Species: *Myotis luficugus*, Little brown myotis

Date surveyed: 6 July 2016

Emergence count 2016: 435 individuals; estimated colony size: 417

 Bat exit/entrance point


 Flight pathway



Figure 4: Little brown myotis (*Myotis luficugus*) maternity roost RM-16-02

Roost size (length x width x height): 11 m x 9 m x 7 m

General location: Mentasta, AK

Comments: The roost is an abandoned house/cabin situated on private land. Roof materials are wood covered with shingle rolls.

Roost ID: RM-16-03

Species: *Myotis luficugus*, Little brown myotis

Date surveyed: 7 July 2016

Emergence count 2016: 57 individuals

 Bat exit/entrance point


 Flight pathway



Figure 5: Little brown myotis (*Myotis luficugus*) maternity roost RM-16-03

Roost size (length x width x height): 4 m x 4 m x 4 m

General location: Nabesna Road, AK

Comments: The roost is a guest cabin used frequently during the summer, situated on private land. Infrequently heated using wood stove. Roof material is tin. Bats roost along the spine of the roof during the day, between the closed ceiling and roofing material. An artificial bat box (style: rocket box) was built during July 2016 and installed approximately 10 meters away on the south side of the building (see appendix B for details).

Roost ID: RM-16-04

Species: *Myotis luficugus*, Little brown myotis

Date surveyed: 10 July 2016

Emergence count 2016: 468 individuals

⦿ Bat exit/entrance point

→ Flight pathway



Figure 6: Little brown myotis (*Myotis luficugus*) maternity roost RM-16-04

Roost size (length x width x height): 6 m x 5 m x 7 m

General location: Gakona, AK

Comments: The roost is a private residence with human occupants year round. The roof material is tin. Bats roost all throughout the roof which majority of the bats congregating along the spine. An artificial bat box (style: rocket box) was installed approximately 50 meters from the roost on the northwest side of the building during July 2016 (see appendix B for details).

Roost ID: RM-16-05

Species: *Myotis luficugus*, Little brown myotis

Date surveyed: 12 July 2016

Emergence count 2016: 83 individuals; estimated colony size: 66

⊙ Bat exit/entrance point

→ Flight pathway



Figure 7: Little brown myotis (*Myotis luficugus*) maternity roost RM-16-05

Roost size (length x width x height): 20 m x 7 m x 8 m

General location: Kenny Lake, AK

Comments: The roost is a private residence with human occupants year round. Numerous attempts have been made to block the roost entrance points. Six artificial bat houses were erected at the roost during autumn 2015: three were attached to the southwest side of the roost building, and three were attached to two outbuildings on the northwest side of the roost. An additional artificial roost (style: rocket box) was installed approximately twenty behinds from the roost, along the treeline and directly on the bat emergence flight path on the south side of the roost building, during July 2016 (see appendix B for details).

Roost ID: RM-16-06

Species: *Myotis luficugus*, Little brown myotis

Date surveyed: 19 July 2016

Emergence count 2016: 84 individuals; estimated colony size: 45

 Bat exit/entrance point


 Flight pathway



Figure 8: Little brown myotis (*Myotis luficugus*) maternity roost RM-16-06

Roost size* (length x width x height): 10 m x 6 m x 9 m

*Bats are limited to the arctic entry: 3 m x 2 m x 3 m

General location: Kenny Lake, AK

Comments: Bats were primarily concentrated in the top portion of the 'arctic entry' on the south side of the building. Historically bats were present throughout the cabin roof/ceiling, however the home owners filled majority of the entrance points which resulted in the bats restricting use to the arctic entry.

APPENDIX B – Artificial Roost Pilot Project

Artificial Roost ID: RB-16-01

Target Species: *Myotis lucifugus*, Little brown myotis

Date installed: 10 July 2016



Figure 9: Installation of artificial bat maternity roost RB-16-01 (style: rocket box) behind an active *Myotis lucifugus* maternity roost (RM-16-03).

General Location: Installed behind RM-16-03. Located on Nabesna Road, AK.

Comments: Installed along tree line behind RM-16-03. Not on major flight path of emerging bats.

Artificial Roost ID: RB-16-02

Target Species: *Myotis luficugus*, Little brown myotis

Date installed: 12 July 2016




Figure 10: Construction of artificial bat roost RB-16-02 (style: rocket box).

General Location: Installed behind RM-16-04. Located in Gakona, AK.

Comments: Installed along tree line, directly on the flight path of bats emerging from RM-16-04.

Artificial Roost ID: RB-16-03

Target Species: *Myotis luficugus*, Little brown myotis

Date installed: 14 July 2016



Figure 11: Installation of artificial bat maternity roost RB-16-03 (style: rocket box) behind an active *Myotis lucifugus* maternity roost (RM-16-05; not in picture).

General Location: Installed behind RM-16-05, near Kenny Lake, AK.

Comments: Installed along tree line, directly on the flight path of bats emerging from RM-16-05.

APPENDIX C – Banding Records - 2016

Table 5: Little brown myotis (*Myotis lucifugus*) banding records for captures during July 2016 surveys

| Band | Date | Location | Species | Sex | Age Class | Researcher |
|--------|----------|----------|---------|-----|-----------|-----------------|
| AP0319 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0320 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0321 | 8-Jul-16 | RM-16-03 | MYLU | M | Adult | J. Reimer, ACCS |
| AP0322 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0323 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0324 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0325 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0326 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0327 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0328 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0329 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0330 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0331 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0332 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0333 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0334 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0335 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0336 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0337 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0338 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0339 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0340 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0341 | 8-Jul-16 | RM-16-03 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0342 | 9-Jul-16 | RM-16-02 | MYLU | F | Juvenile | J. Reimer, ACCS |
| AP0343 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0344 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0345 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0346 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0347 | 4-Jul-16 | RM-16-01 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0348 | 4-Jul-16 | RM-16-01 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0349 | 4-Jul-16 | RM-16-01 | MYLU | M | Adult | J. Reimer, ACCS |
| AP0350 | 4-Jul-16 | RM-16-01 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0351 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0352 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0353 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0354 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0355 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |

| Band | Date | Location | Species | Sex | Age Class | Researcher |
|--------|-----------|----------|---------|-----|-----------|-----------------|
| AP0356 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0357 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0358 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0359 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0360 | 9-Jul-16 | RM-16-02 | MYLU | M | Adult | J. Reimer, ACCS |
| AP0361 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0362 | 9-Jul-16 | RM-16-02 | MYLU | M | Adult | J. Reimer, ACCS |
| AP0363 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0364 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0365 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0366 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0367 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0368 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0369 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0370 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0371 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0372 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0373 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0374 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0375 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0376 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0377 | 9-Jul-16 | RM-16-02 | MYLU | M | Juvenile | J. Reimer, ACCS |
| AP0378 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0379 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0380 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0381 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0382 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0383 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0384 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0385 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0386 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0387 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0388 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0389 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0390 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0391 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0392 | 9-Jul-16 | RM-16-02 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0393 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0394 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0395 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0396 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0397 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |

| Band | Date | Location | Species | Sex | Age Class | Researcher |
|--------|-----------|----------|---------|-----|-----------|-----------------|
| AP0398 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0400 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0401 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0402 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0403 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0404 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0405 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0406 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0407 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0408 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0409 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0410 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0411 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0412 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0413 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0414 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0415 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0416 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0417 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0418 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0419 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0420 | 11-Jul-16 | RM-16-04 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0421 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0422 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0423 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0424 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0425 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0426 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0427 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0428 | 13-Jul-16 | RM-16-05 | MYLU | F | Juvenile | J. Reimer, ACCS |
| AP0429 | 13-Jul-16 | RM-16-05 | MYLU | F | Juvenile | J. Reimer, ACCS |
| AP0430 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0431 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0432 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0433 | 13-Jul-16 | RM-16-05 | MYLU | F | Juvenile | J. Reimer, ACCS |
| AP0434 | 13-Jul-16 | RM-16-05 | MYLU | F | Juvenile | J. Reimer, ACCS |
| AP0435 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0436 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0437 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0438 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0439 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0440 | 13-Jul-16 | RM-16-05 | MYLU | F | Adult | J. Reimer, ACCS |

| Band | Date | Location | Species | Sex | Age Class | Researcher |
|--------|-----------|----------|---------|-----|-----------|-----------------|
| AP0441 | 20-Jul-16 | RM-16-06 | MYLU | F | Juvenile | J. Reimer, ACCS |
| AP0442 | 20-Jul-16 | RM-16-06 | MYLU | F | Juvenile | J. Reimer, ACCS |
| AP0443 | 20-Jul-16 | RM-16-06 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0444 | 20-Jul-16 | RM-16-06 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0445 | 20-Jul-16 | RM-16-06 | MYLU | F | Unknown | J. Reimer, ACCS |
| AP0446 | 20-Jul-16 | RM-16-06 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0447 | 20-Jul-16 | RM-16-06 | MYLU | F | Adult | J. Reimer, ACCS |
| AP0448 | 20-Jul-16 | RM-16-06 | MYLU | F | Juvenile | J. Reimer, ACCS |
| AP0449 | 20-Jul-16 | RM-16-06 | MYLU | F | Juvenile | J. Reimer, ACCS |
| AP0450 | 20-Jul-16 | RM-16-06 | MYLU | F | Juvenile | J. Reimer, ACCS |
| AK1801 | 20-Jul-16 | RM-16-06 | MYLU | F | Adult | J. Reimer, ACCS |
| AK1802 | 20-Jul-16 | RM-16-06 | MYLU | F | Juvenile | J. Reimer, ACCS |
| AK1803 | 20-Jul-16 | RM-16-06 | MYLU | M | Juvenile | J. Reimer, ACCS |
| AK1804 | 20-Jul-16 | RM-16-06 | MYLU | F | Adult | J. Reimer, ACCS |
| AK1805 | 20-Jul-16 | RM-16-06 | MYLU | F | Adult | J. Reimer, ACCS |