

Wilson's Warbler

Cardellina pusilla

Class: Aves
Order: Passeriformes

Conservation Status

NatureServe: Agency:

G Rank: G5 BLM: IUCN: Least Concern Audubon AK: Watch

S Rank: S5B USFWS: ADF&G: Species of Greatest Conservation Need

Final Rank		
Conservation category: V. Orange		
V = unknown status and either high biological vulnerability or high action need		
<u>Category</u>	<u>Range</u>	<u>Score</u>
Status:	-20 to 20	0
Biological:	-50 to 50	-32
Action:	-40 to 40	4
Higher numerical scores denote greater concern		

Status - variables measure the trend in a taxon's population status or distribution. Higher status scores denote taxa with known declining trends. Status scores range from -20 (increasing) to 20 (decreasing).

	Score
<i>Population Trend (-10 to 10)</i>	0
Suspected to be decreasing in interior Alaska (Schmidt et al. 2013; Handel and Sauer 2017). Trends in southeast Alaska appear to be stable (Handel and Sauer 2017). Trends are unknown for other parts of its Alaskan range. We rank this question as Unknown.	
<i>Distribution Trend (-10 to 10)</i>	0
Unknown.	
Status Total:	0

Biological - variables measure aspects of a taxon's distribution, abundance and life history. Higher biological scores suggest greater vulnerability to extirpation. Biological scores range from -50 (least vulnerable) to 50 (most vulnerable).

	Score
<i>Population Size (-10 to 10)</i>	-10
Uncertain, but >25,000. PIF (2019) estimates the population size in Alaska at 35 million individuals (95% CI: 25 to 48 million). Handel et al. (2009) estimated that ~261,000 individuals breed in the Yukon-Charley Rivers National Preserve.	
<i>Range Size (-10 to 10)</i>	-10
Breeds throughout most of the state except for the Arctic coastal plain and the western portion of the Seward Peninsula (Kessel 1989; Ammon and Gilbert 1999). Possible breeder in the eastern Aleutian Islands (Gibson and Byrd 2007). Breeding range is >400,000 sq. km. Overwinters from Mexico south to Panama (Ammon and Gilbert 1999).	
<i>Population Concentration (-10 to 10)</i>	-10
Does not concentrate (Ammon and Gilbert 1999).	
<i>Reproductive Potential</i>	
<u>Age of First Reproduction (-5 to 5)</u>	-5

Few data available, but thought to be 1 year (Ammon and Gilbert 1999).		
<u>Number of Young (-5 to 5)</u>		1
In Alaska, clutch sizes ranging from 3 to 7 have been reported (Ammon and Gilbert 1999), with average clutch sizes of 4.5 ± 0.1 SE (Ammon and Gilbert 1999) and 5.6 ± 0.6 (Kessel 1989). Thought to lay only one clutch per year, though double-brooding has been reported elsewhere in its range (Ammon and Gilbert 1999).		
<i>Ecological Specialization</i>		
<u>Dietary (-5 to 5)</u>		1
Few data available. Consumes adult and larval invertebrates including spiders, beetles, ants, flies, and butterflies (Ammon and Gilbert 1999). In Alaska, Kessel (1989) noted that individuals were also observed eating berries and seeds, though the importance of plant matter to its diet has not been studied.		
<u>Habitat (-5 to 5)</u>		1
Habitat preferences appear to vary regionally in Alaska (Cotter and Andres 2000a). It is most common in tall shrubs and woodland-shrub habitats (Isleib and Kessel 1973; Petersen et al. 1991; Johnson et al. 2008b; Amundson et al. 2018; Savage et al. 2018), but it has also been found in a variety of forest types (e.g. Isleib and Kessel 1973; Matsuoka et al. 2001), including spruce stands with little understory (Petersen et al. 1991). Nevertheless, habitats with thick, tall shrubs appear to support higher densities (Cotter and Andres 2000a; Matsuoka et al. 2001; Amundson et al. 2018).		
Biological Total:		-32
Action - variables measure current state of knowledge or extent of conservation efforts directed toward a given taxon. Higher action scores denote greater information needs due of lack of knowledge or conservation action. Action scores range from -40 (lower needs) to 40 (greater needs).		Score
<hr/> <i>Management Plans and Regulations (-10 to 10)</i>		2
Protected under the Migratory Bird Treaty Act (MBTA 1918).		
<i>Knowledge of Distribution and Habitat (-10 to 10)</i>		-10
Habitat associations and distribution are well-known throughout the state from many multi-species surveys and studies (see Habitat section; Ruthrauff et al. 2007; Schmidt et al. 2013; Corcoran et al. 2014). Some knowledge of migration timing in different parts of its Alaskan range (Benson et al. 2006).		
<i>Knowledge of Population Trends (-10 to 10)</i>		2
Although it is commonly detected during multi-species surveys (see above), population monitoring is only conducted in parts of the interior and southeast Alaska, mostly through BBS and ALMS or localized monitoring programs (e.g. Schmidt et al. 2013). Trends are available from these locations (see Population trends), but statewide trends are not available.		
<i>Knowledge of Factors Limiting Populations (-10 to 10)</i>		10
Little is known about the factors that limit populations in Alaska. A study on Kodiak Island from 2010 to 2014 found that Wilson's warblers exhibited extremely low levels of productivity, both overall and when compared to older data from elsewhere in its Alaskan range (Corcoran et al. 2014). Moreover, both Schmidt et al. (2013) and Handel and Sauer (2017) have documented declines in interior Alaska. However, contributing factors are unknown. A long-term study in central California found that abundance was positively correlated to recruitment in the previous year, and only weakly correlated with high overwinter survival (Chase et al. 1997). The authors concluded that population dynamics in their study were mostly influenced by factors on breeding grounds (Chase et al. 1997). Other studies have found that habitat loss at stop-over and wintering sites negatively affect body condition, but additional research is needed to link these findings to population dynamics (Yong et al. 1998; Ruiz-Sánchez et al. 2017). It is unknown whether these findings apply to populations in Alaska.		
Several studies have investigated the use of genetics to investigate migratory routes and connectivity across its North American range (e.g. Kimura et al. 2002; Clegg et al. 2003; Irwin et al. 2011; Paxton et al. 2013; Ruegg et al. 2014; Kardynal et al. 2018). Breeding populations in interior Alaska appear to be most closely related to populations that breed in Alberta and share similar migratory ecology (Ruegg et al. 2014). Additional research is needed to determine levels of differentiation between eastern and western populations in North America, which		

may represent separate species (Ruiz-Sánchez et al. 2015).

Action Total: 4

Supplemental Information - variables do not receive numerical scores. Instead, they that are used to sort taxa to answer specific biological or management questions.

Harvest:	None or Prohibited
Seasonal Occurrence:	Breeding
Taxonomic Significance:	Monotypic species
% Global Range in Alaska:	>10%
% Global Population in Alaska:	25-74%
Peripheral:	No

References

- Ammon, E. M. and W. M. Gilbert. 1999. Wilson's Warbler (*Cardellina pusilla*), version 2.0. In Poole, A. F., and F. B. Gill, eds. *The Birds of North America*, Cornell Lab of Ornithology, Ithaca, NY, USA. DOI: 10.2173/bna.478
- Amundson, C. L., C. M. Handel, D. R. Ruthrauff, T. L. Tibbitts, and R. E. Gill. 2018. Montane-breeding bird distribution and abundance across national parks of southwestern Alaska. *Journal of Fish and Wildlife Management* 9(1):180–207. DOI: 10.3996/062017-
- Benson, A. M., B. A. Andres, W. N. Johnson, S. Savage, and S. M. Sharbaugh. 2006. Differential timing of Wilson's warbler migration in Alaska. *The Wilson Journal of Ornithology* 118(4):547–551. DOI: 10.1676/05-085.1
- Chase, M. K., N. Nur, and G. R. Geupel. 1997. Survival, productivity, and abundance in a Wilson's warbler population. *The Auk* 114(3):354–366. DOI: 10.2307/4089238
- Clegg, S. M., J. F. Kelly, M. Kimura, and T. B. Smith. 2003. Combining genetic markers and stable isotopes to reveal population connectivity and migration patterns in a Neotropical migrant, Wilson's warbler (*Wilsonia pusilla*). *Molecular Ecology* 12(4):819–
- Corcoran, R., C. Trussell, and R. MacIntosh. 2014. Monitoring Avian Productivity and Survivorship on Kodiak Island, Alaska, 2010-2014. Refugue report 2014.7, Kodiak National Wildlife Refuge, U.S. Fish and Wildlife Service, Kodiak, AK, USA.
- Cotter, P. A., and B. A. Andres. 2000a. Breeding bird habitat associations on the Alaska breeding bird survey. Information and Technology Report USGS/BRD/ITR- 2000-0010, Biological Resource Division, U.S. Geological Survey, Springfield, VA, USA.
- Gibson, D. D., and G. V. Byrd. 2007. *Birds of the Aleutian Islands, Alaska*. Nuttall Ornithological Club, Cambridge, MA, USA.
- Handel, C. M. and Sauer, J. R. 2017. Combined analysis of roadside and off-road breeding bird survey data to assess population change in Alaska. *The Condor* 119(3):557-575. DOI: 10.1650/CONDOR-17-67.1
- Handel, C. M., S. A. Swanson, D. A. Nigro, and S. M. Matsuoka. 2009. Estimation of avian population sizes and species richness across a boreal landscape in Alaska. *The Wilson Journal of Ornithology* 121(3):528–547.
- Irwin, D. E., J. H. Irwin, and T. B. Smith. 2011. Genetic variation and seasonal migratory connectivity in Wilson's warblers (*Wilsonia pusilla*): species-level differences in nuclear DNA between western and eastern populations. *Molecular Ecology* 20(15):310
- Isleib, M. E., and B. Kessel. 1973. *Birds of the north Gulf Coast- Prince William Sound region, Alaska*. Biological Papers of the University of Alaska no. 14. University of Alaska Fairbanks, AK, USA.

- Johnson, J. A., B. A. Andres, and J. A. Bissonette. 2008b. Birds of the major mainland rivers of Southeast Alaska. General Technical Report PNW-GTR-739. U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, Portland, OR, USA.
- Kardynal, K. J., D. M. Collister, and K. A. Hobson. 2018. Origins of Wilson's warblers migrating through southwest Canada: adding value to banding data by using stable isotopes and genetic markers. *Animal Migration* 5(1):17–28. DOI: 10.1515/ami-2018-0002
- Kessel, B. 1989. Birds of the Seward Peninsula, Alaska: Their biogeography, seasonality, and natural history. University of Alaska Press, Fairbanks, AK, USA.
- Kimura, M., S. M. Clegg, I. J. Lovette, K. R. Holder, D. J. Girman, B. Mila, P. Wade, and T. B. Smith. 2002. Phylogeographical approaches to assessing demographic connectivity between breeding and overwintering regions in a Nearctic-Neotropical warbler (W)
- Matsuoka, S. M., C. M. Handel, and D. R. Ruthrauff. 2001. Densities of breeding birds and changes in vegetation in an Alaskan boreal forest following a massive disturbance by spruce beetles. *Canadian Journal of Zoology* 79(9):1678–1690. DOI: 10.1139/cjz-79
- Migratory Bird Treaty Act (MBTA). 1918. U.S. Code Title 16 §§ 703-712 Migratory Bird Treaty Act.
- Paxton, K. L., M. Yau, F. R. Moore, and D. E. Irwin. 2013. Differential migratory timing of western populations of Wilson's warbler (*Cardellina pusilla*) revealed by mitochondrial DNA and stable isotopes. *The Auk* 130(4):689–698. DOI: 10.1525/auk.2013.13107
- Petersen, M. R., D. N. Weir, and M. H. Dick. 1991. Birds of the Kilbuck and Ahklun Mountain Region, Alaska. *North American Fauna* 76:1-158.
- Partners in Flight (PIF). 2019. Population Estimates Database, version 3.0. Available online: <http://pif.birdconservancy.org/PopEstimates>. Accessed 09-April-2019.
- Ruegg, K. C., E. C. Anderson, K. L. Paxton, V. Apkenas, S. Lao, R. B. Siegel, ..., and T. B. Smith. 2014. Mapping migration in a songbird using high-resolution genetic markers. *Molecular Ecology* 23(23):5726–5739. DOI: 10.1111/mec.12977
- Ruiz-Sánchez, A., K. Renton, R. Landgrave-Ramírez, E. F. Mora-Aguilar, and O. Rojas-Soto. 2015. Ecological niche variation in the Wilson's warbler *Cardellina pusilla* complex. *Journal of Avian Biology* 46(5):516–527. DOI: 10.1111/jav.00531
- Ruiz-Sánchez, A., K. Renton, and R. Rueda-Hernández. 2017. Winter habitat disturbance influences density and territory size of a Neotropical migratory warbler. *Journal of Ornithology* 158(1):63–73. DOI: 10.1007/s10336-016-1368-9
- Ruthrauff, D. R., T. L. Tibbitts, R. E. Gill, and C. M. Handel. 2007. Inventory of montane-nesting birds in Katmai and Lake Clark National Parks and Preserves. Report NPS/AKRSWAN/NRTR-2007/02, U.S. Geological Survey Alaska Science Center, Anchorage, AK, U
- Savage, S., T. L. Tibbitts, K. Sesser, and R. S. A. Kaler. 2018. Inventory of lowland-breeding birds on the Alaska Peninsula. *Journal of Fish and Wildlife Management*. In press. DOI: 10.3996/082017-JFWM-070
- Schmidt, J. H., C. L. McIntyre, and M. C. MacCluskie. 2013. Accounting for incomplete detection: what are we estimating and how might it affect long-term passerine monitoring programs? *Biological Conservation* 160:130–139. DOI: 10.1016/j.biocon.2013.01.007
- Yong, W., D. M. Finch, F. R. Moore, and J. F. Kelly. 1998. Stopover ecology and habitat use of migratory Wilson's warblers. *The Auk* 115(4):829–842. DOI: 10.2307/4089502

Review status: Peer-reviewed

Version date: 21 May 2019

Alaska Center for Conservation Science
Alaska Natural Heritage Program
University of Alaska Anchorage
Anchorage, AK