

Dark-eyed Junco (oreganus)

Class: Aves
Order: Passeriformes

Junco hyemalis oregonus

Note: This assessment refers to this subspecies only. A species level report, which refers to all associated subspecies, is also available.

Review Status: Peer-reviewed

Version Date: 09 May 2019

Conservation Status

NatureServe: Agency:

G Rank: G5T5 ADF&G: Species of Greatest Conservation Need IUCN: Audubon AK:

S Rank: USFWS: BLM:

Final Rank		
Conservation category: IX. Blue		
low status and low biological vulnerability and action need		
<u>Category</u>	<u>Range</u>	<u>Score</u>
Status	-20 to 20	-6
Biological	-50 to 50	-20
Action	-40 to 40	0
Higher numerical scores denote greater concern		

Status - variables measure the trend in a taxon's population status or distribution. Higher status scores denote taxa with known declining trends. Status scores range from -20 (increasing) to 20 (decreasing).

Score

Population Trend in Alaska (-10 to 10)

-6

Short-term (2003-2015) trends are stable or increasing in southcoastal and southeast Alaska (Handel and Sauer 2017). Long-term trends (1993-2015) are stable (Handel and Sauer 2017). Potentially declining elsewhere in its range in the Pacific Northwest (Pardieck et al. 2017; Rockwell et al. 2017).

Distribution Trend in Alaska (-10 to 10)

0

Unknown.

Status Total: -6

Biological - variables measure aspects of a taxon's distribution, abundance and life history. Higher biological scores suggest greater vulnerability to extirpation. Biological scores range from -50 (least vulnerable) to 50 (most vulnerable).

Score

Population Size in Alaska (-10 to 10)

0

Unknown.

Range Size in Alaska (-10 to 10)

-2

Breeds in southeast Alaska north to Yakutat Bay (Gabrielson and Lincoln 1959; Andres et al. 2005). A portion of the breeding population overwinters in southcoastal and southeast Alaska (Nolan et al. 2002; Armstrong 2008), while the rest overwinters further south (Nolan et al. 2002). Breeding range

is most restricted and is estimated to be ~88,090 sq. km, based on range map from ACCS (2017a).

Population Concentration in Alaska (-10 to 10) -10

Does not concentrate (Nolan et al. 2002).

Reproductive Potential in Alaska

Age of First Reproduction (-5 to 5) -5

Breeds within its first year (Nolan et al. 2002).

Number of Young (-5 to 5) 1

Unknown for Alaska. Elsewhere in North American, clutch size ranges from 3 to 5 per clutch (Nolan et al. 2002). In northern parts of their range, lays only one clutch per year (Nolan et al. 2002).

Ecological Specialization in Alaska

Dietary (-5 to 5) -5

Unknown for Alaska. Elsewhere in North America, juncos are omnivorous and consume mainly seeds and invertebrates including spiders, wasps, ants, and beetles (Nolan et al. 2002). The percent of vegetable versus plant matter in their diet appears to change seasonally with availability (Nolan et al. 2002).

Habitat (-5 to 5) 1

Nests on the ground and forages in a variety of forest types, stand ages, and disturbance regimes (Dellasala et al. 1996; Lance and Howell 2000; Cotter and Andres 2000a; Matsuoka et al. 2001). Prefers open canopy forests and tends to avoid areas with a thick shrub understory (Matsuoka et al. 2001; Matsuoka and Handel 2007); however, this species is occasionally detected in tall shrub habitat (Spindler and Kessel 1980; Kessler and Kogut 1985; Cotter and Andres 2000a).

Biological Total: -20

Action - variables measure current state of knowledge or extent of conservation efforts directed toward a given taxon. Higher action scores denote greater information needs due of lack of knowledge or conservation action. Action scores range from -40 (lower needs) to 40 (greater needs).

Score

Management Plans and Regulations in Alaska (-10 to 10) 2

Protected under the Migratory Bird Treaty Act (MBTA 1918).

Knowledge of Distribution and Habitat in Alaska (-10 to 10) -10

Detected during multi-species surveys in southeast and southcoastal Alaska, with knowledge of habitat associations (e.g. Kessler and Kogut 1985; Dellasala et al. 1996; Willson and Gende 2000; Johnson et al. 2008b).

Knowledge of Population Trends in Alaska (-10 to 10) -2

Commonly detected during multi-species surveys in appropriate habitat (Cotter and Andres 2000a). Trend information is available from Breeding Bird Surveys and off-road surveys conducted in southeast and southcoastal Alaska (Handel and Sauer 2017).

Knowledge of Factors Limiting Populations in Alaska (-10 to 10) 10

Few studies have considered the population dynamics of *J. hyemalis* in Alaska or elsewhere. Some factors have been proposed, including nest predation and weather, but limiting factors have not been identified. In southcentral Alaska, Matsuoka and Handel (2007) found that predation was the main cause of nest failure. Predation was mediated by spruce bark beetle infestations. Nest success was lowest in forest stands that were least affected by spruce bark beetle (Matsuoka and Handel 2007).

They attributed this difference to the higher rates of nest predation by red squirrels, which are closely associated with intact spruce forests (Matsuoka et al. 2001; Matsuoka and Handel 2007). Spruce bark beetle and other disturbances such as logging may also increase local abundances of dark-eyed juncos, which prefer open canopy forests (Dellasala et al. 1996; Lance and Howell 2000). Willson and Gende (2000) reported high rates of nesting success in southeast Alaska, but did not identify factors that may influence reproductive success. Inclement weather on overwintering grounds can lead to annual fluctuations in population size (reviewed in Nolan et al. 2002). In Alaska, climate change may affect timing of arrival on breeding grounds (Mizel et al. 2017) and may increase suitable habitat if the treeline moves further north or higher up, as predicted by climate models (Marcot et al. 2015; Mizel et al. 2016). Studies elsewhere in this species' range have reported geographic differences in reproductive parameters across elevational gradients, potentially resulting from differences in the length of the breeding season and in food availability (Bears et al. 2009; LaBarbera and Lacey 2018). In Alberta's Rocky Mountains, dark-eyed juncos breeding at high-elevations had lower reproductive success, but produced higher-quality offspring (Bears et al. 2009). Timing of snowmelt, which also exhibits an elevational gradient, may also affect clutch size and phenology (Smith and Andersen 1985; DeSante 1990). It is unknown whether similar differences in reproductive parameters exist across latitudinal gradients, though Nolan et al. (2002) documented that double-brooding only occurs in southern parts of their global range.

Action Total: 0

Supplemental Information - variables do not receive numerical scores. Instead, they are used to sort taxa to answer specific biological or management questions.

Harvest:	None or Prohibited
Seasonal Occurrence:	Year-round
Taxonomic Significance:	Subspecies
% Global Range in Alaska:	>10%
% Global Population in Alaska:	25-74%
Peripheral:	No

References

- Alaska Center for Conservation Science (ACCS). 2017a. Wildlife Data Portal. University of Alaska Anchorage. Available online: <http://aknhp.uaa.alaska.edu/apps/wildlife>
- Andres, B. A., B. T. Browne, and D. L. Brann. 2005. Composition, abundance, and timing of post-breeding migrant landbirds at Yakutat, Alaska. *The Wilson Bulletin* 117(3):270–279. DOI: 10.1676/04-039.1
- Armstrong, R. H. 2008. Guide to the birds of Alaska, 5th edition. Alaska Northwest Books, Anchorage, AK, USA.
- Bears, H., K. Martin, and G. C. White. 2009. Breeding in high-elevation habitat results in shift to slower life-history strategy within a single species. *Journal of Animal Ecology* 78(2):365–375. DOI: 10.1111/j.1365-2656.2008.01491.x
- Cotter, P. A., and B. A. Andres. 2000a. Breeding bird habitat associations on the Alaska breeding bird survey. Information and Technology Report USGS/BRD/ITR- 2000-0010, Biological Resource Division, U.S. Geological Survey, Springfield, VA, USA.
- Dellasala, D. A., J. C. Hagar, K. A. Engel, W. C. McComb, R. L. Fairbanks, and E. G. Campbell. 1996. Effects of silvicultural modifications of temperate rainforest on breeding and wintering bird communities, Prince of Wales Island, Southeast Alaska. *The Condor* 98(4):706–721. DOI: 10.2307/1369853
- DeSante, D. F. 1990. The role of recruitment in the dynamics of a Sierran subalpine bird community. *The American Naturalist* 136(4):429-445.

- Gabrielson, I. N., and F. C. Lincoln. 1959. *The Birds of Alaska*. The Stackpole Company, Harrisburg, PA, USA.
- Handel, C. M. and Sauer, J. R. 2017. Combined analysis of roadside and off-road breeding bird survey data to assess population change in Alaska. *The Condor* 119(3):557-575. DOI: 10.1650/CONDOR-17-67.1
- Johnson, J. A., B. A. Andres, and J. A. Bissonette. 2008b. *Birds of the major mainland rivers of Southeast Alaska*. General Technical Report PNW-GTR-739. U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, Portland, OR, USA.
- Kessler, W. B., and T. E. Kogut. 1985. Habitat orientations of forest birds in southeastern Alaska. *Northwest Science* 59(1):58-65.
- LaBarbera, K., and E. A. Lacey. 2018. Breeding season length and nest mortality drive cryptic life history variation in Dark-eyed Juncos (*Junco hyemalis*) breeding across a montane elevational gradient. *The Auk* 135(2):284–298. DOI: 10.1642/AUK-17-184.1
- Lance, E. W., and S. Howell. 2000. Survey of songbirds during a spruce beetle (*Dendroctonus rufipennis*) outbreak on the Kenai Peninsula, Alaska. *Northwestern Naturalist* 81(1):1-10. DOI: 10.2307/3536893.
- Marcot, B. G., M. T. Jorgenson, J. P. Lawler, C. M. Handel, and A. R. DeGange. 2015. Projected changes in wildlife habitats in Arctic natural areas of northwest Alaska. *Climate Change* 130(2):145–154. DOI: 10.1007/s10584-015-1354-x
- Matsuoka, S. M., and C. M. Handel. 2007. Nesting ecology of boreal forest birds following a massive outbreak of spruce beetles. *Journal of Wildlife Management* 71(1):51–63. DOI: 10.2193/2005-460
- Matsuoka, S. M., C. M. Handel, and D. R. Ruthrauff. 2001. Densities of breeding birds and changes in vegetation in an Alaskan boreal forest following a massive disturbance by spruce beetles. *Canadian Journal of Zoology* 79(9):1678–1690. DOI: 10.1139/cjz-79-9-1678
- Migratory Bird Treaty Act (MBTA). 1918. U.S. Code Title 16 §§ 703-712 Migratory Bird Treaty Act.
- Mizel, J. D., J. H. Schmidt, C. L. McIntyre, and C. A. Roland. 2016. Rapidly shifting elevational distributions of passerine species parallel vegetation change in the subarctic. *Ecosphere* 7(3):e01264. DOI: 10.1002/ecs2.1264
- Mizel, J. D., J. H. Schmidt, C. L. McIntyre, and M. S. Lindberg. 2017. Subarctic-breeding passerines exhibit phenological resilience to extreme spring conditions. *Ecosphere* 8(2):e01680. DOI: 10.1002/ecs2.1680
- Nolan Jr., V., E. D. Ketterson, D. A. Cristol, C. M. Rogers, E. D. Clotfelter, ..., and E. Snajdr. 2002. Dark-eyed Junco (*Junco hyemalis*), version 2.0. In Poole, A. F., and F. B. Gill, eds. *The Birds of North America*, Cornell Lab of Ornithology, Ithaca, NY, USA. DOI: 10.2173/bna.716
- Pardieck, K. L., D. J. Ziolkowski Jr., M. Lutmerding, K. Campbell and M.-A.R. Hudson. 2017. North American Breeding Bird Survey Dataset 1966 - 2016, version 2016.0. U.S. Geological Survey, Patuxent Wildlife Research Center. DOI:10.5066/F7W0944J.
- Rockwell, S. M., J. D. Alexander, J. L. Stephens, R. I. Frey, and C. J. Ralph. 2017. Spatial variation in songbird demographic trends from a regional network of banding stations in the Pacific Northwest. *The Condor* 119(4):732–744. DOI: 10.1650/CONDOR-17-44.1
- Smith, K. G., and D. C. Andersen. 1985. Snowpack and variation in reproductive ecology of a montane ground-nesting passerine, *Junco hyemalis*. *Scandinavian Journal of Ornithology* 16(1):8–13.
- Spindler, M. A., and B. Kessel. 1980. Avian populations and habitat use in interior Alaska taiga. Final report, University of Alaska Museum, Fairbanks, AK, USA.
- Willson, M. F., and S. M. Gende. 2000. Nesting success of forest birds in Southeast Alaska and adjacent Canada. *The Condor* 102:314–325.