

# White-winged Scoter

*Melanitta fusca*

Class: Aves  
Order: Anseriformes

**Review Status:** Peer-reviewed

**Version Date:** 21 March 2018

## Conservation Status

NatureServe: Agency:

G Rank: G5      ADF&G: Species of Greatest Conservation Need      IUCN: Least Concern      Audubon AK:

S Rank: S5B,S5N      USFWS:      BLM:

<b>Final Rank</b>		
Conservation category: <b>VII. Yellow</b>		
low status and either high biological vulnerability or high action need		
<u>Category</u>	<u>Range</u>	<u>Score</u>
Status	-20 to 20	-3
Biological	-50 to 50	-28
Action	-40 to 40	4
<b>Higher numerical scores denote greater concern</b>		

**Status** - variables measure the trend in a taxon’s population status or distribution. Higher status scores denote taxa with known declining trends. Status scores range from -20 (increasing) to 20 (decreasing).

**Score**

*Population Trend in Alaska (-10 to 10)*

2

It is difficult to detect trends for this species because it is hard to differentiate between white-winged scoters and other scoters. Consequently, scoter species are often lumped together during the main survey for breeding waterfowl in North America (the USFWS Waterfowl Breeding Population and Habitat Survey). Moreover, the timing of this survey is inadequate for late-season nesters such as scoters and scaup. Nevertheless, some data are available. Surveys in Alaska for the three scoter species that breed in the state suggest declines from the 1980s to the 2000s and increasing trends since then (Flint 2013; Bowman et al. 2015). Species-specific data indicate an increasing trend in Alaska between 1993 and 2012 (Bowman et al. 2015), but the authors still consider their population size to be below historic levels. Population estimated on the Yukon Flats National Wildlife Refuge (NWR), which supports the largest population of white-winged scoters in the state, have been stable for the past 13 years (Guldager et al. 2016).

*Distribution Trend in Alaska (-10 to 10)*

-5

Although no formal studies have been conducted, its distribution is suspected to be stable (T. Lewis, ADF&G, pers. comm.). There is no evidence of contractions to their breeding or wintering distribution and the habitats they occupy (winter - temperate marine, breeding - boreal forest) are thought to be stable (T. Lewis, ADF&G, pers. comm.).

Status Total: -3

**Biological** - variables measure aspects of a taxon's distribution, abundance and life history. Higher biological scores suggest greater vulnerability to extirpation. Biological scores range from -50 (least vulnerable) to 50 (most vulnerable).

**Score***Population Size in Alaska (-10 to 10)*

-10

Unknown, but likely >25,000. Surveys on the Yukon Flats NWR, which support the largest breeding population of white-winged scoters in the state (qtd. in. Guldager et al. 2016), estimate a population index of 15,403 individuals (based on a 13-year mean; Guldager et al. 2016).

*Range Size in Alaska (-10 to 10)*

-8

Breeds in boreal forest wetlands across interior Alaska from the Canadian border west to Innoko and Koyukuk, south to the Wrangell Mountains, and north to the Brooks Range and the Arctic National Wildlife Refuge (Johnson and Herter 1989; ACCS 2017a; Brown and Fredrickson 2019). Summer sightings in southeast Alaska (Hodges et al. 2008) and on the Arctic Coastal Plain (Johnson and Herter 1989) are of non-breeding and post-breeding individuals, respectively. Overwinters from the Aleutian Islands east to the Gulf of Alaska and along the coast to southeast Alaska. Wintering range is most restricted and is estimated at ~200,000 sq. km., estimated in GIS.

*Population Concentration in Alaska (-10 to 10)*

-10

White-winged scoters typically migrate in small flocks up to 100 birds (Brown and Fredrickson 2019), though larger groups of several hundred individuals have been reported in winter and during molting (e.g. Isleib and Kessel 1973; Butler 1998; Heint and Piston 2009). Coastal and offshore waters from Prince William Sound to southeast Alaska are important areas during migration (Isleib and Kessel 1973; Butler 1998; USFWS 2011) and overwinter (Hodges 2011; USFWS 2011). Important staging areas have also been identified in southwest Alaska e.g. Port Moller and Ugashik Bay (USFWS 2011). A large number of (presumed) non-breeding birds also remain in southeast Alaska during the summer (Hodges et al. 2008). Given population size and its widespread distribution year-round, number of sites is likely >250.

*Reproductive Potential in Alaska*Age of First Reproduction (-5 to 5)

-3

Breeds at two years of age (Brown and Fredrickson 2019).

Number of Young (-5 to 5)

1

Average clutch size is between 8.3 to 9.6 eggs and ranges from 5 to 19 (Brown and Brown 1981; Brown and Fredrickson 2019). On the Yukon Flats, mean clutch size was  $8.1 \pm 0.2$  eggs (Safine 2005). However, a large proportion of the adult female population (28%) did not breed every year (Safine 2005). Females lay a single clutch per year and do not reneest in the event of nest failure (Brown and Fredrickson 2019; T. Lewis, ADF&G, pers. comm.).

*Ecological Specialization in Alaska*Dietary (-5 to 5)

1

Consumes mainly benthic freshwater invertebrates on breeding grounds (crustaceans, insect larvae) and benthic marine invertebrates on wintering grounds (Anderson et al. 2008; Brown and Fredrickson 2019). Scoters' winter dietary niche has been described as "generalists on molluscs, mostly bivalves" (Sanger and Jones 1984). Indeed, bivalves are an especially important prey item in winter (Sanger and Jones 1984; Anderson et al. 2008; Palm et al. 2012). When bivalves are unavailable, white-winged scoters are flexible in their dietary choices, consuming fish, crustaceans, polychaete worms, and echinoderms (Palm et al. 2012). Scoters are also capable of responding to pulsed resources e.g. altering movement and foraging behaviors in response to herring spawning events (Lewis et al. 2007; Lok et al. 2008; Heint and Piston 2009).

Habitat (-5 to 5)

1

Breeds in boreal and prairie-parkland habitats. Tends to avoid graminoid sites, preferring to place

their nest in shrub or forested habitat that provide good nesting cover or escape from predators (e.g. on islands) (Brown and Brown 1981; Safine and Lindberg 2008). While nests are often located far from shore (Brown and Brown 1981; Safine and Lindberg 2008), Safine (2005) considered White-winged scoters to be generalist within the broader shrub-forest habitat, placing nests at various distances from the shoreline. Young are reared on lakes in shallow, vegetated areas (Brown and Brown 1981; Safine 2005). On the Yukon Flats, occupancy of lakes by brood-rearing scoters was best explained by lake size >100ha and high amphipod densities (Lewis et al. 2015). Overwinters in protected coastal areas such as bays and inlets that support abundant mollusc densities (Lewis et al. 2008b; Gunn 2009; Brown and Frederickson 2019).

Biological Total: -28

**Action** - variables measure current state of knowledge or extent of conservation efforts directed toward a given taxon. Higher action scores denote greater information needs due of lack of knowledge or conservation action. Action scores range from -40 (lower needs) to 40 (greater needs).

**Score**

*Management Plans and Regulations in Alaska (-10 to 10)*

-10

Protected by the Migratory Bird Treaty Act (MBTA 1918). Subsistence harvest and recreational hunting is permitted, but subject to regulations (ADFG 2018e; AMBCC 2018). An action plan is in place for this species (USFWS 2011).

*Knowledge of Distribution and Habitat in Alaska (-10 to 10)*

2

Little is known about distribution and habitat use in Alaska, though recent work has furthered our understanding. Studies have addressed migratory movements, identified staging and molting areas, and characterized winter habitat requirements (Gunn 2009; reviewed in USFWS 2011). Safine (2005) was the first to study the breeding ecology of white-winged scoters in Alaska. More recently, Lewis et al. (2015) investigated habitat selection of brood-rearing scoters. Both of these studies were conducted on the Yukon Flats NWR. Our knowledge of other breeding populations in Alaska remains scarce.

*Knowledge of Population Trends in Alaska (-10 to 10)*

2

Current surveys are inadequate for detecting trends. Because it is hard to differentiate between scoter species, scoters are often lumped together during surveys (Bowman et al. 2015). Moreover, the main survey for breeding waterfowl in North America (the USFWS Waterfowl Breeding Population and Habitat Survey) is inadequately timed for late-season nesters such as scoters and scaup (Guldager et al. 2016). To address these limitations, USFWS has flown annual scoter and scaup surveys on the Yukon Flats NWR since 2001 (Guldager et al. 2016). Other localized surveys occur sporadically and are reviewed in USFWS (2011).

*Knowledge of Factors Limiting Populations in Alaska (-10 to 10)*

10

White-winged scoters are difficult to study during the breeding season because of their propensity for nesting in low densities across vast landscapes. Little is known about their ecology and the factors that limit its population in Alaska or elsewhere (USFWS 2011). Compared to studies in Canada, researchers on the Yukon Flats noted lower nest and female survival, but higher duckling survival (Safine 2005). Nearly 30% of adult females did not breed every year, which may act to further restrict population growth rates (Safine 2005). Predation is the main cause of mortality for eggs, ducklings, and females on breeding grounds (Safine 2005; Brown and Fredrickson 2019). Although white-winged scoters rely heavily on bivalves in the winter, food does not appear to be limiting in at least some parts of their range (Lewis et al. 2008b). The scoter's large body size may offer some protection from variations in food resources and energetic stressors during winter and molting seasons (Anderson and Lovvorn 2011; Uher-Koch 2013). It is unknown whether hunting is a significant source of mortality (Safine 2005). Subsistence harvest data are limited but can exceed 12,500 birds per year in Alaska (Naves 2015).

Large die-offs have been reported on molting grounds in southeast Alaska and in Washington (Henny 1995; USFWS 2011; Jones et al. 2017). In southeast Alaska, the cause for the die-off was unknown: no infectious diseases were found, and levels of some contaminants were high but largely inconclusive (Henny 1995). In Washington, the die-off was correlated with the timing of a harmful algal bloom (HAB; Jones et al. 2017). HABs may become an increasingly serious threat for marine birds as the frequency and magnitude of these blooms have been linked to climate change (Jones et al. 2017).

Action Total: 4

**Supplemental Information** - variables do not receive numerical scores. Instead, they are used to sort taxa to answer specific biological or management questions.

<b>Harvest:</b>	Substantial, regulations
<b>Seasonal Occurrence:</b>	Year-round
<b>Taxonomic Significance:</b>	Monotypic species
<b>% Global Range in Alaska:</b>	>10%
<b>% Global Population in Alaska:</b>	<25%
<b>Peripheral:</b>	No

## References

- Alaska Center for Conservation Science (ACCS). 2017a. Wildlife Data Portal. University of Alaska Anchorage. Available online: <http://aknhp.uaa.alaska.edu/apps/wildlife>
- Alaska Department of Fish and Game (ADFG). 2020c. 2020-2021 Migratory game bird hunting regulations summary. Anchorage, AK, USA.
- Anderson, E. M., and J. R. Lovvorn. 2011. Contrasts in energy status and marine foraging strategies of white-winged scoters (*Melanitta fusca*) and surf scoters (*M. perspicillata*). *The Auk* 128(2):248-257. DOI: 10.1525/auk.2011.10088.
- Anderson, E. M., J. R. Lovvorn, and M. T. Wilson. 2008. Reevaluating marine diets of surf and white-winged scoters: Interspecific differences and the importance of soft-bodied prey. *The Condor* 110(2):285-295. DOI: 10.1525/cond.2008.8458.
- Bowman, T. D., E. D. Silverman, S. G. Gilliland, and J. B. Leirness. 2015. Status and trends of North American sea ducks: Reinforcing the need for better monitoring. Pages 1-28 in J.-P. L. Savard, D. V. Derksen, D. Esler, and J. M. Eadie, eds. *Ecology and conservation of North American sea ducks*. Studies in Avian Biology No. 46. CRC Press, Boca Raton, FL, USA.
- Brown, P. W., and M. A. Brown. 1981. Nesting biology of the white-winged scoter. *Journal of Wildlife Management* 45(1):38-45. DOI: 10.2307/3807871.
- Brown, P. W., and L. H. Fredrickson 2019. White-winged Scoter (*Melanitta deglandi*), version 1.1. In Rodewald, P. G., ed. *The Birds of North America*, Cornell Lab of Ornithology, Ithaca, NY, USA. DOI: 10.2173/bna.whwsco4.01.1
- Butler, R. W. 1998. Moulting sites of sea ducks and other marine birds in Frederick Sound, Southeast Alaska. *Canadian Field-Naturalist* 112(2):346-347.
- Flint, P. L. 2013. Changes in size and trends of North American sea duck populations associated with North Pacific oceanic regime shifts. *Marine Biology* 160(1):59-65. DOI: 10.1007/s00227-012-2062-y
- Guldager, N., M. Bertram, and B. Lake. 2016. 2014 and 2015 aerial scoter and scaup surveys, Alaska. Page 22. Yukon Flats National Wildlife Refuge Report – 2016-001, U.S. Fish and Wildlife Service, Yukon Flats National Wildlife Refuge, Fairbanks, AK, USA.
- Gunn, T. 2009. Habitat correlates of wintering sea duck occurrence in Southeast Alaska. MSc thesis, Simon Fraser University, Burnaby, BC, CAN.

- Heinl, S. C., and A. W. Piston. 2009. Birds of the Ketchikan area, Southeast Alaska. *Western Birds* 40(2):54–144.
- Henny, C. J., D. D. Rudis, T. J. Roffe, and E. Robinson-Wilson. 1995. Contaminants and sea ducks in Alaska and the circumpolar region. *Environmental Health Perspectives* 103(Suppl. 4):41-49.
- Hodges, J. I. 2011. Exploratory winter sea duck survey of southcentral Alaska – Cape Spencer to Prince William Sound. Available online: <http://seaduckjv.org/pdf/studies/pr129.pdf>
- Hodges, J. I., D. J. Groves, and B. P. Conant. 2008. Distribution and abundance of waterbirds near shore in Southeast Alaska, 1997-2002. *Northwestern Naturalist* 89(2):85-96.
- Isleib, M. E., and B. Kessel. 1973. Birds of the north Gulf Coast- Prince William Sound region, Alaska. *Biological Papers of the University of Alaska* no. 14. University of Alaska Fairbanks, AK, USA.
- Johnson, S. R., and D. R. Herter. 1989. The birds of the Beaufort Sea. BP Exploration Inc., Anchorage, AK, USA.
- Jones, T., J. K. Parrish, A. E. Punt, V. L. Trainer, R. Kudela, J. Lang, ..., and B. Hickey. 2017. Mass mortality of marine birds in the Northeast Pacific caused by *Akashiwo sanguinea*. *Marine Ecology Progress Series* 579:111-127. DOI: 10.3354/meps12253
- Lewis, T. L., D. Esler, and W. S. Boyd. 2007. Foraging behaviors of surf scoters and white-winged scoters during spawning of Pacific herring. *The Condor* 109(1):216-222.
- Lewis, T. L., D. Esler, and W. S. Boyd. 2008b. Foraging behavior of surf scoters (*Melanitta perspicillata*) and white-winged scoters (*M. fusca*) in relation to clam density: Inferring food availability and habitat quality. *The Auk* 125(1):149-157. DOI: 10.1525/auk.2008.125.1.149.
- Lewis, T. L., M. S. Lindberg, J. A. Schmutz, M. R. Bertram, and A. J. Dubour. 2015. Species richness and distributions of boreal waterbird broods in relation to nesting and brood-rearing habitats. *Journal of Wildlife Management* 79(2):296-310. DOI: 10.1002/jwmg.837
- Lok, E. K., M. Kirk, D. Esler, and W. S. Bond. 2008. Movements of pre-migratory surf and white-winged scoters in response to Pacific herring spawn. *Waterbirds* 31(3):385-393.
- Migratory Bird Treaty Act (MBTA). 1918. U.S. Code Title 16 §§ 703-712 Migratory Bird Treaty Act.
- Naves, L. C. 2015. Alaska subsistence bird harvest, 2004-2014 data book. Special Publication No. 2015-05, Alaska Department of Fish and Game, Division of Subsistence, Anchorage, AK, USA.
- Palm, E. C., D. Esler, E. M. Anderson, and M. T. Wilson. 2012. Geographic and temporal variation in diet of wintering white-winged scoters. *Waterbirds* 35(4):577-589.
- Safine, D. E. 2005. Breeding ecology of white-winged scoters on the Yukon Flats, Alaska. MSc thesis, University of Alaska Fairbanks, AK, USA.
- Safine, D. E., and M. S. Lindberg. 2008. Nest habitat selection of white-winged scoters on Yukon Flats, Alaska. *Wilson Journal of Ornithology* 120(3):582-593.
- Sanger, G. A., and R. D. Jones. 1984. The winter feeding ecology and trophic relationships of oldsquaws and white-winged scoters on Kachemak Bay, Alaska. In Nettleship, D. N., G. A. Sanger, and P. R. Springer, eds. *Marine birds: their feeding ecology and commercial fisheries relationships*. Proceedings of the Pacific Seabird Group Symposium, 6-8 January 1982, Seattle, WA, USA.
- Uher-Koch, B. 2013. Latitudinal and seasonal variation in non-breeding survival of surf and white-winged scoters. MSc thesis, Simon Fraser University, Burnaby, BC, CAN.
- U.S. Fish and Wildlife Service (USFWS). 2011. Action plan for white-winged scoter. Unpublished report, U.S. Fish and Wildlife Service, Anchorage, AK, USA. Available online: <https://www.fws.gov/migratorybirds/pdf/management/focal-species/White-wingedScoter.pdf>

